
Recommendations For Management Of Children With Diabetes In School







Recommendations for Management of Children with Diabetes in School was developed under a grant from the Nevada State Diabetes Control Program.

Nevada Diabetes Association for Children and Adults 1005 Terminal Way, Suite 104 Reno, NV 89502

> 775.856.3839 ndaca@diabetesnv.org http://diabetesnv.org

TABLE OF CONTENTS

AKNOWLEDGEMENTS	3
INTRODUCTION	4
REGULATIONS AFFECTING SCHOOL POPULATIONS	5
ABOUT DIABETES	7
CONTROL OF DIABETES	8
CARE PLANNING	10
PARENT CONFERENCE	11
PLANNING MEETING	11
INDIVIDUAL CARE PLAN	12
TRAINING	13
DAILY MANAGEMENT	14
BLOOD GLUCOSE MONITORING PROCEDURES	15
HYPOGLYCEMIA (LOW BLOOD GLUCOSE)	18
HYPOGLYCEMIA (LOW BLOOD GLUCOSE)—WITH LOSS OF CONSCIOUSNESS OR SEIZURE	19
HYPERGLYCEMIA (HIGH BLOOD GLUCOSE)	21
KETOACIDOSIS	21
FOOD AND SPECIAL OCCASIONS	23
EXERCISE AND SPORTS	26
NSULIN	27
ROLES AND RESPONSIBILITIES	28
APPENDIX 1	36
COMPARISON OF MANIFESTATIONS OF HYPOGLYCEMIA AND HYPERGLYCEMIA	36
APPENDIX 2	37
AUTHORIZATION FOR RELEASE OF MEDICAL INFORMATION	37
APPENDIX 3	38

PRESCRIPTION MEDICATION ORDER AND PERMISSION FORM	38
APPENDIX 4	39
MEDICAL STATEMENT FOR CHILDREN	39
REQUIRING MODIFICATIONS IN SCHOOL MEALS	39
APPENDIX 5	40
SAMPLE504 PLAN FOR THE SCHOOL YEAR 97/98	40
APPENDIX 6	43
SAMPLE STUDENT ACCOMMODATION PLAN	43
UNIVERSAL PRECAUTIONS AND INFECTION CONTROL	44
INSTRUCTIONS FOR COMPLETING THE MEDICAL STATEMENT	46
DATA/INFORMATION SHEET—DIABETES CARE IN SCHOOL	47
PARENT CHECKLIST	51
INDIVIDUAL CARE PLAN— DIABETES CARE IN SCHOOL	52
SUPPLEMENTAL INFORMATION SHEET FOR STUDENT WEARING AN INSULIN PUMP AT SCHOOL (LICENCED HEATHCARE PROVIDER'S ORDERS FOR INSULIN PUMPS FOR HEATH SERVICES AT SCHOOL)	

AKNOWLEDGEMENTS

The Nevada Diabetes Association for Children and Adults developed this manual under a grant from the Nevada State Diabetes Control Program. The manual is based on 'Recommendations for Management of Diabetes for Children in School', developed by Eileen Crawford, BSN, RN for the Vermont Department of Health, Diabetes Control Program.

The following individuals contributed to the creation of the Vermont manual: Jo Busha, Nathaniel Clark, MD, Paul Costello, MD, Paula Duncan, MD, Nancy Emberley, Barbara Farrell, RD, Donna Filipek, Erin Filipek, Fiona Gagnon, Mary Alice Giannoni, NP, Mary Gill, RN, PNP, Carlie Geer, RD, MS, Bonnie Hudspeth, Lynn Hurley, RN, CDE, Sue Johansen, RD, CDE, Linda LaShure, RD, CD, Marilyn McSweeney, RN, Amy Nickerson, RD, Alison Precourt, RD, CDE, Pam Spinney, Margaret Terrien, MSN, CDE, Elizabeth Warren, Janice Whitaker, RN.

We would also like to thank Jeff Hitchcock of Children with Diabetes http://www.childrenwithdiabetes.com/index_cwd.htm for assembling the information on the School Bill of Rights for Children with Diabetes and the web sites for additional information.

'Recommendations for Management of Children with Diabetes in School' was edited from the original by Mylan Hawkins, Executive Director, NDACA with the assistance of Kathy Eckert, M.D., Carolyn Leontos, MS, RD, and Kim Neiman, Control Program Manager and the suggestions and guidance of the members of the Nevada Diabetes Council.

INTRODUCTION

This manual has been prepared to assist school staff to assure that children, who have diabetes, and their families, have the support needed to manage the disease in the school setting. The Nevada State Diabetes Control Program (DCP) has been established to reduce the burden of diabetes for people with diabetes and their families.

The Nevada Diabetes Council (NDC) is a coalition of individuals, organizations, businesses and corporations concerned with diabetes that advises the DPC regarding its program and assists the DPC in areas of Public and Professional Education, Access to Care and Public Policy, Data Collection and information and referrals to resources throughout the state and nation.

One compelling need identified through the NDC is the management and safety of children with diabetes in school. Parents throughout the state have identified many instances where schools have not responded appropriately to the needs and the rights of their children. Examples include: not allowing children with diabetes to go on field trips or to participate in sporting events; denying modification of school meals when needed; interference with how the child tests their blood glucose and/or manages their insulin therapy, and unwillingness to train staff to assist children to manage their disease in the school setting.

Specific goals of the manual include:

- To assure the safety of the child is maintained in the school setting through development and implementation of an individual care plan and training of school staff.
- To support the student in becoming independent with their self-care management that is consistent with their age and interest.
- To guide schools in their understanding of diabetes and its management; and, assure the coordination of diabetes care in school with the care provided in the home.
- To enhance opportunities for children with diabetes to participate fully in all school functions.
- The manual is designed to be copied and distributed to school staff, parents and others as deemed appropriate and is available on diskette if requested. It may also be seen and printed from the Internet at the following addresses. http://health2k.state.nv.us/diabetes/ or http://diabetesnv.org

REGULATIONS AFFECTING SCHOOL POPULATIONS

School Bill of Rights for Children with Diabetes

Children with diabetes require medical care to remain healthy. The need for medical care does not end while the child is at school. Thus, while at school, each child with diabetes must be allowed to:

- 1. Do blood sugar checks
- 2. Treat hypoglycemia with emergency sugar
- 3. Inject insulin when necessary
- 4. Eat snacks when necessary
- 5. Eat lunch at an appropriate time and have enough time to finish the meal
- 6. Have free and unrestricted access to water and the bathroom Participate fully in physical education (gym class) and other extracurricular activities, including field trips

The Laws

The right of children with diabetes to care for their diabetes at school is based on the **Individuals With Disability Education Act** (IDEA) and **Section 504** of the Rehabilitation Act of 1973. These laws provide protection against discrimination for children with disabilities, including diabetes. Parents can use these laws to ensure that, while at school, their children with diabetes can fully participate in all school activities, while at the same time caring for their medical needs.

Any school that receives Federal funding must comply with IDEA and Section 504 laws. A child need not require special education to be protected.

Taking Action Against Discrimination

If you are faced with a school that does not comply with the **School Bill of Rights for Children with Diabetes**, you should first educate the school administration. Make sure they understand the laws and your child's needs.

Schools that still refuse to cooperate should be advised that you are requesting preparation of an **Individualized Education Program** (IEP) and a **Section 504 Accommodation** for your child. At this point, the school must meet with you to negotiate the special services that your child requires. You should begin with the entire list of services in the **School Bill of Rights for Children with Diabetes**. If your school still refuses to comply with the **School Bill of Rights for Children with Diabetes**, you should file a complaint with your state's department of education. This is the first step in the process of litigation against your school system.

Your child has a right to care for his or her diabetes at school. The scientific data are clear on the value of maintaining glycemic control. Since there is no break from diabetes, there can be no break from the need to care for it. Time spent at school is no exception.

For Additional Information

Your School and Your Rights, http://www.diabetes.org/ada/scrights.asp from the American Diabetes Association, discusses the legal obligations of school systems under Section 504 of the Rehabilitation Act of 1973 and the Education for All Handicapped Children Act of 1975, amended in 1991.

The National Information Center for Children and Youth with Disabilities (NICHCY) http://www.nichcy.org/ is a U.S. Government sponsored clearinghouse that provides information about disabilities, including information about obtaining assistance at school. Important information on their web site includes:

The *Individuals with Disabilities Education Act (IDEA) Amendments of 1997*. President Clinton signed the bill into law on June 4, 1997.

State Contact Sheets, with the names and contact information for state officials and education offices to which parents can turn for assistance.

The US Department of Education http://www.ed.gov/ web site includes:
The *Individuals with Disabilities Education Act*, with detailed information about IDEA. *IDEA: The Law* contains links to downloadable versions of the law.

Education Administration Online http://phoenix.lrp.com/EAO/ publishes a newsletter on *Inclusive Education Programs*.

The **Department of Justice ADA Home Page**

http://www.usdoj.gov/crt/ada/adahom1.htm has references to the law and related links. The DOJ ADA site also includes the *KinderCare Settlement Agreement re: Diabetes Finger-Prick Tests*, which includes information about diabetes for use by KinderCare employees.

Section 504 from the U.S. Department of Labor. **How to File a Complaint with the Office of Civil Rights** http://www.hhs.gov/ocr/hivfile.html under Section 504, courtesy of the U.S. Department of Health and Human Services.

How to Write an I.E.P. http://www.childrenwithdiabetes.com/iep is a book designed to help parents who have children with disabilities succeed in school. Many parents of children with diabetes use an IEP to ensure that their children can care for their diabetes in school.

ABOUT DIABETES

Description

Diabetes is a chronic metabolic disorder, resulting in the body's inability to utilize glucose for energy due to a lack of or a resistance to insulin. Diabetes is not contagious. The cause is unknown, but appears to be a combination of genetic and environmental factors.


Type 1 diabetes, previously called juvenile, or insulin dependent diabetes, is an autoimmune disease in which the insulin producing cells of the pancreas are destroyed, resulting in the need for daily injections of insulin. Insulin is necessary for blood glucose (also called "blood sugar") to be maintained at normal levels.

Type 2 diabetes, previously called adult onset, or non-insulin dependent diabetes, was considered a rare occurrence in children but is becoming more common, especially in adolescents. Children who develop type 2 diabetes are overweight and usually have a strong family history of diabetes. A resistance to insulin rather than a deficiency of insulin characterizes type 2 diabetes.

The information in this manual applies to both type 1 diabetes and type 2 diabetes.

CONTROL OF DIABETES

There is no cure for diabetes. The treatment plan is directed at managing or controlling its course. Diabetes is controlled by balancing insulin or oral medication, food and exercise. Blood glucose testing enables people with diabetes to monitor these three components of management and adjust one or more of them to correct or maintain a good balance.



Low blood sugar, *hypoglycemia*, may be caused by too little food, too much insulin or more exercise than usual. Elevated blood sugar, *hyperglycemia*, may be caused by too much food, too little insulin, illness, stress or less exercise than usual.

The child's school performance will be best if the blood glucose remains in the acceptable range. Left untreated both high and low blood glucose levels can effect the child's ability to concentrate on schoolwork and participate in school activities.

Appendix 1 provides a comparison of hypo and hyperglycemia.

Psychosocial Concerns

Trust. Parents need to be reassured that the school will be able to manage their child's diabetes related needs. Working closely with the parents in planning for the child's care, by training all staff, and by establishing clear lines of communication and responsibility among team members will enhance trust.

Developmental level. It is important to consider the child with diabetes within the context of their developmental level, both currently and at the age of diagnosis. Developmental level will determine, to a large degree, the child's acceptance and response to the disease and level of participation in self-care. It is important that the parent(s)/guardian provide the school information on the level of assistance their child will need. (See Forms Data/Information Sheet—page 47) It is equally important that all those who interface with the child in the school environment understand what to do if called on to assist the child and be sensitive to the fact that even a child that is self-sufficient in managing their care may at times need assistance.

Living with chronic illness. A condition, such as diabetes, that interferes with daily functioning for more than three months in a year is defined as a chronic illness. Like all chronic conditions, diabetes has a profound affect on the family. Planning for the care and meeting the needs of a child with diabetes can be exhausting. Family coping mechanisms should be supported and enhanced so that acute and chronic complications for the child can be avoided.

The child with diabetes is subject to many feelings about his/her condition. School personnel can be instrumental in assisting the child and his/her family to accept the disease and be prepared to cope with behaviors related to denial, fear, anger, guilt and depression.

Self-empowerment. The child living with diabetes must learn to integrate the many facets of self-care diabetes management into their individual life-style. Children may be assisted along this continuum with the careful consideration of individual capabilities. Providing the opportunity to develop skills in self-care will promote self-confidence. Ultimately, a child's improved self-concept will go far toward achieving better control over the disease.

Adolescence. Many adolescent issues stem from the need to "fit in" or not appear to be different from peers. Fearing rejection, the adolescent may attempt to hide the fact that he/she has diabetes. In some instances, denial may lead to a deliberate rejection of components of self-management, which may result in poor disease control. Other issues that may interfere with the successful management of diabetes in adolescents include weight awareness and/or body image, particularly with females, as well as depression, which are related to self-image or disease management.

The school team should be sensitive to the adolescent who is struggling with issues related to their diabetes. In some situations, referral to counseling may be appropriate. Good communication with parents and other influential adults may assist passage through this tumultuous developmental stage. School personnel can offer important support and guidance for families throughout adolescence to foster the positive growth and development of adolescents living with diabetes.

CARE PLANNING

Care planning in the school has four components:

Parent Conference. A conference with parents, and the school nurse to identify the child's needs, discuss components of the care plan, and develop the agenda for a school wide planning meeting.

Planning Meeting. This meeting of key staff should be held in the fall before the school year starts, or when a child is newly diagnosed.

"Individual Care Plan". The School Nurse, using information gathered at the planning meeting, should prepare the written plan. Key staff and the child's family must agree to the plan. The plan may be incorporated into a "504" plan if the child's needs will be covered by this legislation. See Regulations (page 4) and Guidelines (page 5) for a description of legislative rules that may apply to children with diabetes.

Training. The school nurse should arrange for training for all school staff. The nurse should do the training with the assistance of the child's parents and/or invited members of the child's health care team. This may involve one or more sessions depending on the roles assigned to different people.

NOTE: The Nevada Diabetes Resource Directory contains resources that may help the school staff with care planning. This directory is available on and can be downloaded from the internet, at

http://health2k.state.nv.us/diabetes/Diabetes%20Council/index.htm or http://icanv.com/ndaca It is also available in hard copy and on disk from the Diabetes Control Program at (775) 684-5949 or (775) 856-3839.

Parent Conference

This meeting is held with parents, the principal, the school nurse and others who may be invited by one of these parties. The purpose is to get to know one another, share information about the child and school, prepare for the initial planning meeting, and determine who will need to attend the planning meeting.

The parent checklist should be prepared at the parent conference so that the necessary forms and supplies can be brought to the planning meeting.

Sample forms that the school might want to use are included in this Guide. The information needed from the parent is included on the suggested form. This may be completed by the parent in advance of the parent conference and brought with them or filled out during the conference.

Planning Meeting

Each fall, and at other times during the school year for children who are newly diagnosed, the school nurse in the school district should organize and facilitate a planning meeting to develop an individual diabetes care plan for use in the school setting.

Meeting participants should include everyone that may have a role in the child's diabetes care. Participants may include:

- Family and child
- Principal
- School nurse
- Current year classroom teacher(s)
- Past year classroom teacher(s)
- Food service manager
- Physical education teacher/coach
- Counselor or Social Worker
- Bus driver
- Other school staff with direct responsibility for child
- Members of the health care team, if invited by parents

Suggested agenda items:

- Overview of Type 1 or Type 2 Diabetes and its management
- Roles and responsibilities of staff members
- Identify staff in the school who will serve as resources for others
- Determine the hierarchy of personnel expected to respond to emergency situations

- Determine the location of food kits, glucagon and other supplies in the school building
- Determine where the plan will be kept and how individual components will be shared with appropriate staff
- How training for staff with specific responsibilities will be done
- What is an emergency and what to do

Individual Care Plan

Planning is key to the successful management of care for children with diabetes. In schools, the individual care plan is a good tool for accomplishing successful management.

The school nurse, in collaboration with parents and others, develops this care plan outlining specific health care to be given to the student. It should be available to all staff working with the child. The school nurse may extract sections of the plan or copy pages from this manual clip them to the care plan and distribute them to each team member.

Routine daily care includes:

- Blood glucose monitoring routine
- Phone numbers of parents, guardians, care providers and emergency contacts
- Blood glucose values and specify responses required
- Daily schedule of food, insulin and activity
- Special events/circumstances
- Location of supplies and food
- Disposal of syringes, lancets, etc.
- Developmental levels and cognitive and physical abilities of the school-aged child and adolescent should be incorporated in the care plan for the child with diabetes

Emergencies:

Children with diabetes can have problems despite the best efforts at control. School staff needs to determine and record what constitutes an emergency situation and what to do about it. Parents and the health care team should provide guidance for the care plan.

Sample Care Plans:

A suggested sample care plan that can be done by a school nurse is included in this Guide along with other forms. These have integrated aspects of the individual care plan that follows with a School 504 Plan and a Student Accommodation Plan.

Training

Goals for training: Everyone mentioned in the plan knows their role in carrying out the plan, how it relates to the roles of others and when and where to seek help.

Preparation: Assess school personnel to determine their knowledge of and comfort level with caring for the student. Modify the training session accordingly.

Time: The initial session should take about 30 minutes. Some members of the staff may need additional individual training around their specific roles.

Attendance: Include all staff mentioned in the plan plus administrative and counseling staff and any others who may interact with the child during the school session.

Suggested components of training:

- Introduction to the child's individual care plan
- Type 1 Diabetes or Type 2 Diabetes: what it is, how it is managed (if not covered at planning meeting)
- Monitoring tools: glucometer, ketone kit, written records, etc.
- Signs and symptoms of hypoglycemia and hyperglycemia
- Procedures for routine care of the individual student
- Emergency procedures
- Overview of universal health and safety guidelines (OSHA) and disposal of supplies.
- Monitoring techniques (for those who may do finger sticks)
- Glucagon administration (for those named in the emergency plan)
- Insulin administration (if in the plan)

DAILY MANAGEMENT

Blood glucose monitoring is done to check the child's blood glucose level so that appropriate decisions about the balance of insulin, food and exercise can be made.

This section describes how to perform monitoring as well as how to interpret and act upon the results.

Monitoring Procedures: Times, record keeping and specific procedures are established at the planning meeting and included in the individual care plan.

Hypoglycemia: Low blood glucose is the most common event that will need attention in school. Generally, it is not an emergency situation. Hypoglycemia is discussed in detail in the section that follows.

Hypoglycemia with loss of consciousness: An *emergency* response is indicated, guidance is included.

Hyperglycemia: High blood glucose may also be seen in school. It is also described in some detail with suggested interventions presented. Hyperglycemia is generally not an emergency situation.

Ketoacidosis: The result of relatively long-term elevation in blood glucose levels, ketoacidosis requires action, but will rarely constitute an emergency.

Parents should inform the schools as to what levels require specific intervention for their child in order to avoid either high or low blood glucose levels. This information should be in the individual care plan.

BLOOD GLUCOSE MONITORING PROCEDURES

Checking blood glucose:

Children with both Type 1 and Type 2 diabetes may need to check their blood glucose at school. If the child needs assistance with blood glucose monitoring the person assisting the child should report the specific values to the child and describe blood glucose results as "high, low, or normal". Do not use terms like "good or bad," attitudes are generally improved when feelings of good or bad, right or wrong are not attached.

Check with parents for guidance as to which term you should use. Keep them consistent with those used by the child's family and primary health care team.

Who:

The school nurse, teacher or other staff member who has been trained should do the monitoring for children needing assistance. Older children will usually prefer to carry their own monitoring/insulin kits and do their own checking.

When:

Glucose monitoring in school should closely follow the monitoring schedule done at home. In general, monitoring for children with type 1 diabetes is done four times per day, usually before meals and before bedtime. In addition, some children may need monitoring before or after exercise, and all children will need additional tests if they exhibit symptoms of high or low blood glucose or are ill. The care plan should specify when regular monitoring is to occur and when symptoms require.

Where:

Monitoring should be done wherever a child with diabetes feels most comfortable in the school setting and where it will cause the least disruption to the child's academic program. This may be in the classroom if that is acceptable to the child.

If the child is exhibiting signs of hypoglycemia either accompany the child to the health office or have a designated member of the school team come to the child. This is necessary because the child may become confused and may not be able to make it to the test site alone.

DO NOT ASK another child to be responsible for accompanying a child who may be exhibiting signs of hypoglycemia. Junior or Senior high school students who have been specifically trained and are willing and able to accept the responsibility may be considered **ONLY** if the students and both families agree.

Tools and steps:

Checking the blood glucose level is simple to do. It requires pricking the finger with a lancet to obtain a drop of blood. This is placed on a specially prepared strip. The results are read by a hand held machine (meter) or if a meter is not available, by visual comparison with a color chart located on the bottle. The information is then recorded. The specific steps are shown in the box below.

Consult the student's individualized care plan for specific responses and information pertaining to blood glucose levels, equipment and supply storage and disposal.

Steps for Blood Glucose Monitoring

- 1. Gather supplies.
- 2. Wash hands in warm water.
- 3. Load lancet device with lancet
- 4. Wipe finger with warm soapy water. Let dry. Alcohol swabs should be used only if warm water is not available.
- 5. Hold lancet devise to side of fingertip and press button to stick finger.
- 6. Turn finger down to get a full drop of blood.
- 7. If a larger drop of blood is needed, milk finger.
- 8. Put drop of blood on strip pad.
- 9. Follow directions for using a monitor or to read the result on the bottle of strips.
- 10. Record on sheets provided by parents.

Disposal of supplies:

Cotton balls, swabs or strips with blood: The small amount of blood, with its exposure to air, makes the risk of contamination extremely low. In the home and in the clinic setting disposal is generally done in the regular trash. Schools may prefer to put them in a plastic bag prior to disposal.

Sharps:

Needles and lancets should never be put in the trash. A sharps disposal device should be used if available. If not, a plastic soda bottle with a screw top can be used. The container should be **labeled** and the lid should be taped shut before disposal in the trash.

Record keeping:

Record keeping is fundamental to the optimal management of diabetes. The diabetes record will travel with the child from home to school thereby enabling the communication of patterns of blood glucose values, food intake, variations to normal exercise and the possibility of developing an illness.

Frequent episodes of high or low blood glucose levels in school should alert the school nurse, classroom teacher and others of a need for more communication with the family and/or health care team in order to assess and plan appropriate interventions.

Examples of blood glucose records can be found in resource section of the Appendix.

Classification of numbers:

70-110 mg/dl Normal fasting serum glucose

80-180 mg/dl Goal range for children with diabetes

<70 mg/dl Hypoglycemia >240 mg/dl Hyperglycemia

The individual care plan is the source of appropriate responses to blood glucose values outside of the child's normal range.

Emergency care:

Most situations are easily managed and do not constitute true emergencies. Low blood glucose is the condition most likely to escalate into an emergency in the school setting. Its critical to act quickly at the first signs of hypoglycemia so that an emergency situation does not occur.

An emergency response is required under the following conditions:

- 1. Loss of consciousness, seizure or is unable to take food by mouth. See section on hypoglycemia for appropriate action.
- 2. Blood glucose level does not increase with food ingestion. *Follow parent instructions or section on hypoglycemia.*
- 3. Hypoglycemia with vomiting and with medium or large ketone levels. See section on ketoacidosis.

Hypoglycemia (low blood glucose)

APPROXIMATE LEVEL: < 70 mg/dl

CAUSES: When the body gets too much insulin, meals or snacks are missed or with increased physical activity or exercise.

ONSET: Rapid. Most likely to occur at peak insulin action times, such as before lunch.

SYMPTOMS: Low blood glucose feels different to different people; it may include any of the following. It is important to be aware of a child's specific symptoms.

faintness fatigue shakiness dizziness muscle cramping weakness hunger pale skin

nervousness inappropriate actions

stomachache confusion blurred vision irritability headache crankiness fatigue convulsions

sweating unconsciousness

TREATMENT:

- 1. Test blood glucose. If the child is incoherent, is having difficulty following instructions or is combative, go to step 2.
- Stop activity in order to prevent further reduction in the blood glucose level.
- 3. Eat 15 grams of quick acting carbohydrate glucose such as a small juice box or 4-glucose tablet. If child is having difficulty eating or drinking, Glucose Gel (tubes) may be used. Avoid carbohydrate foods that also contain protein or fat (e.g. chocolate or cookies). These foods are digested more slowly than pure carbohydrate foods.
- 4. Wait 15 minutes.
- 5. Repeat step 1. If glucose level is <80 have the child eat a snack or go to lunch.

NOTE: Over-treatment can result in a rebound effect resulting in high blood glucose levels.

Hypoglycemia (low blood glucose)—WITH Loss of Consciousness or Seizure

EMERGENCY RESPONSE IS INDICATED

Early recognition and treatment of hypoglycemia is extremely important. Left untreated, hypoglycemia can result in loss of consciousness and/or seizure activity. If that occurs, the child may need treatment with the administration of Glucagon, a hormone that causes a rise in blood sugar.

Parents should indicate whether they want the school to administer the Glucagon or to call the Rescue Squad. If the parents wish, they should provide the Glucagon Emergency Kit and the order to administer it (ONLY THE KIT FORM SHOULD BE USED AT SCHOOL). The school should have a list of designated-trained individuals who can administer Glucagon. This list should be kept in an accessible, agreed upon location.

If the administration of Glucagon is not authorized or if there is no one at the school who is properly trained to administer Glucagon the following procedure should be instituted if the child is unconscious or having a seizure.

- 1. If only **one person** is present call the Rescue Squad (911) administer Glucose Gel by squeezing gel into the check. Notify parents.
- 2. If **two people** are present: One should first call the Rescue Squad (911) and then notify the parents, while the other administers Glucose Gel.

Instructions for using Glucagon:

Only a person who has been trained in the following procedure should administer Glucagon. Use *only* when the child is unconscious or having a seizure.

If **two people** are present: One should first call the Rescue Squad (911) and then notify the parents, while the other prepares the Glucagon. If only **one person** is present: First, call 911, then administer the Glucagon and then notify the parents.

- 1. Use the fluid-filled syringe in the Emergency Kit. Inject it into the bottle containing the Glucagon tablet. Shake well to mix.
- 2. Draw Glucagon mix back into the syringe. If the child weighs less than 45 pounds, use ½ of the dose.
- 3. Inject Glucagon intramuscularly into the buttocks or thigh.
- 4. Turn the person onto one side or stomach (vomiting generally occurs with Glucagon).

- 5. After giving the injection give glucose gel by rubbing the glucose gel on the gums and inner cheeks but not actually dumping in the mouth
- 6. Wait 10 minutes after giving Glucagon and administering glucose gel and then check the patient's blood glucose.
- 7. When the patient is awake and drinking, continue giving "quick sugars". Sips of juice, regular (sugared) soda, or sugar in water initially until blood sugar is at least 120. If the child is unable to drink, Glucose gel, honey, or maple syrup may be given to help raise the blood sugar. DO NOT GIVE FOOD THAT CONTAINS FAT as this will slow down the absorption of other sugars. (It may take as much as 20 minutes for the patients blood glucose level to begin to return to normal as generally Glucagon takes about 15-20 minutes to being action.)
- 8. After blood sugar is stable and the patient is not throwing up THEN give solid foods.
- Complete recovery may take 1-2 hours. The effects of Glucagon lasts for 12-25 minutes.

Any child who has been administered Glucagon or has had a seizure should be evaluated and monitored at a hospital until it is clear that the patient is stable and able to keep down fluids and keep the blood sugar up.

Hyperglycemia (high blood glucose)

APPROXIMATE LEVEL: >240 mg/dl

CAUSE: Too little insulin, too much food or too little physical activity. Infections, illness, and/or stress may precipitate the condition.

ONSET: Slow.

SYMPTOMS:

Thirst headaches frequent urination fatigue

dry mouth abdominal pain

ketones in the urine.

TREATMENT:

Check blood glucose levels.

- 2. If blood glucose levels are over 240 mg/dl and/or symptoms of stress or illness are present call parents. Administer extra insulin only if advised by parents.
- 3. Push fluids.
- 4. The child should return to the classroom and their regular routine. They must be allowed to have a water bottle in class and to leave to use the restroom when needed.

NOTE: Exercise should be avoided if ketones are present. Increased physical activity can cause blood glucose to become more elevated and result in further ketone development.

KETOACIDOSIS

Hyperglycemic episodes may result in a serious condition called DKA or diabetic ketoacidosis. When insulin levels are deficient, the cells of the body are starved for energy and will burn fat in response to their need. Ketones are acids produced by the abnormal metabolism of fats.

The risk of DKA occurring at school is extremely low.

ONSET: Can be rapid and can lead to severe illness or even death.

SYMPTOMS:

dehydration

drowsiness

abdominal pain

vomiting

labored breathing and fruity smelling breath.

Adolescents may be at higher risk of DKA than other school age groups.

TREATMENT: (Test as for Hyperglycemia):

- 1. Check blood glucose levels.
- 2. If over 350 and the child has *any* symptoms of ketoacidosis, *call parents immediately*. Administer extra insulin only if advised by parents. Start pushing fluids.
- 3. If the child is vomiting and is unable to take fluids, call the Rescue Squad (911).
- 4. Avoid all forms of exercise.

To Check for Ketones

- 1. Dip ketone test strips in a urine filled paper cup.
- 2. Wait several seconds.
- 3. Read results by visual comparison toa color chart. Refer to instructions printed on the test strip vial.
- 4. Record result for parents.
- 5. Ketone test strips out-date. Be sureto read information on the bottle and discard when out-dated.

FOOD AND SPECIAL OCCASIONS

Food. Planning for meals, snacks, special events and emergency situations for a child with diabetes is an integral part of the diabetes management plan. Each child should have an established meal plan which the child and family has developed with a Registered Dietitian. This meal plan includes time, type and amount of food needed to balance the child's nutritional needs with his/her activity level and insulin regime.

There may be times when the balance of timing of insulin and food with activity and other aspects of the child's life will not be as precise as desired and the child will experience episodes of hypoglycemia. Emergency food supplies, usually called snack packs or low packs by families, will be needed.

Special Occasions. There are **NO** forbidden foods or activities in the lives of people with diabetes. Advance planning will allow the child to participate fully in non-routine activities of the school.

Meal Planning

Food. Meals for children with diabetes are based on the same principles of nutrition as for any child. The primary difference is that the carbohydrate intake for the child with diabetes is usually "controlled" or "planned." Carbohydrates are the body's main source of energy and have the greatest effect on blood glucose levels. They are found in fruit, vegetables, grains, milk and many other foods including those containing sugar. Meats and fats contain no carbohydrate.

The key to good blood glucose control is balancing and spacing carbohydrate food throughout the day at meals and snacks. The **total amount** of carbohydrate eaten at one time is the primary concern, not the **source** of carbohydrate. Research has shown that sugars and sweets included in meals do not raise the blood glucose any quicker than the same amount of carbohydrate from starches or "complex carbohydrates." The healthy nutrition advice suitable for everyone should be followed: "Eat sugar and sweets in moderation."

Snacks. It is often necessary for a child with diabetes to have a snack mid way between breakfast and lunch, depending on his/her insulin regime and activity level. A mid-afternoon snack may be eaten at school or at home depending on the child's individual needs. The care plan should define the timing of snacks and alternatives in case of unforeseen circumstances.

School Meals. The child with diabetes may not need modifications in the school meal. If some changes are necessary, the food service director should be a member of the care plan team and guide the necessary adaptations to the regular menu items. For additional information on school meal modification consult, "Special Meals for Special Needs: A Manual for School Food Service Managers," available from the Department of Education, Child Nutrition Programs.

Federal regulations require that schools participating in the "National School Meals Program" modify meals for children whose disability restricts their diet. A physician must certify the necessary modifications. A copy of the "Medical Statement for Children Requiring Modifications in School Meals" form is included as Appendix.

Emergency Food Supplies ("Snack Packs" or "Low packs"). Good overall planning and access to carbohydrates ensures that the child with diabetes has the means of obtaining appropriate emergency responses during the school day.

Recommended Foods for "Low-Packs"

Food	Grams of
	Carbohydrate
Immediate Treatment	
4 Glucose tablets	16
1 Tube Glucose gel or cake frosting	15
½ cup of Juice	
(orange, apple, grape, pineapple)	15
½ cup of non-diet soda	15
½ cup of Kool-Aid	15
Follow-up Snack (15-30 minutes follow	wing hypoglycemia)
1 cup milk	12
Cracker snack pack(cheese or	
peanut butter)	(as listed on food label)
Granola Bar	(as listed on food label)

The family should furnish emergency food supplies to provide the child's preferred choice of food to respond to a low blood glucose. The food supplies should be in several locations and travel with the child. Appropriate locations for "low packs" might be the health office, physical education office, classroom, school office and school bus.

Special Occasions

Food carries many emotional overtones. It is wise to avoid giving food any special power. This can be accomplished by avoiding terms such as "diet" and "cheat" in dealing with food issues. *Remember* that there are no forbidden foods in a diabetic meal plan.

School Parties. High sugar treats are often high in fat as well and should be discouraged from inclusion at school parties. Providing more nutritious foods, such as low-fat fruit desserts, at school parties is healthier for all children. And serving more nutritious foods gives the message that healthy foods are fun and taste good. This can also serve as a reminder that parties provide an opportunity to share and celebrate important events in our lives; the focus need not be food.

When parties include high sugar/high fat foods notice should be sent to the parents. With appropriate planning children with diabetes may have small amounts of high sugar-high fat party foods, such as cake or ice cream. If no alternative foods are available, the child should be given the same food as everyone else.

Field Trips. Field trips and bus trips also require advance planning. Parents should provide written instructions concerning the special needs of the child while traveling. Supplies that should travel with the child include the items needed to monitor blood glucose level, snack packs, glucose tablets and gel and insulin and/or Glucagon if requested y the parent. The chaperone on the bus and/or the driver, must be informed and provided with instructions regarding the child's care. The child should be allowed to eat on the bus if necessary.

Staying after School. Staying after school should be planned for in advance. Monitoring, mid-afternoon snacks or insulin injections may be part of the routine for children with diabetes and must be accommodated in any after-school activity. This after-school routine should be outlined in the care plan. If the school nurse is not available during after-school events, another staff member should be identified as the primary contact for questions or concerns about the child.

EXERCISE AND SPORTS

In addition to insulin, food and regular monitoring of blood glucose, exercise is a key component of diabetes management. Children and adolescents can fully participate in gym class, team sports, and general playground and individual activities. Professional athletes with diabetes have successfully achieved their physical and athletic potential.

With few special considerations, the student with diabetes can fully participate in the athletic opportunities available in schools. **Program that encourages children to exercise and participate in sports in non-competitive ways can be of special benefit.** Since exercise is an important part of good management of the student's blood glucose level, it should be encouraged on a consistent basis. Children with diabetes can and should be encouraged to "do it all."

Routine Exercise

Exercise improves insulin sensitivity; the duration and intensity of exercise will influence blood glucose levels. This may result in a decreased requirement for insulin and/or an increased potential of hypoglycemic episodes. To avoid this, the student may need to eat an additional snack before exercising. If a child has symptoms of low blood glucose or will be participating in more than 40-45 minutes of physical activity, blood glucose levels should be checked before exercising. Glucose monitoring equipment and a "low pack" should be available at the activity site.

Team Sports

Team sports should be encouraged if the child expresses an interest. The child, parents and care plan will provide the guidance necessary to accommodate full participation. The only accommodations needed will be to have monitoring supplies and snacks available at the site of the activity and staff trained regarding their use. Most children old enough to participate in school sports are able to monitor their own blood and to adjust their snacks accordingly. Many children will chose to sip Gatorade or other sports beverages during practice and games to keep their blood glucose levels high. Children should be allowed this independence.

INSULIN

Children with Type 1 diabetes must inject insulin daily. The physician and family determine the insulin types, dosages and times needed. Insulin needs vary with each child. Most children will not need to inject insulin during the school day. Children who need to inject insulin may need assistance and supervision.

Administration

If the child requires assistance then the school nurse or another designated individual will be responsible for injecting insulin or in assisting a student to draw up and self inject insulin. The procedure for administration of insulin is in the box at the right.

Inspect the insulin. Check the expiration date Printed on the label. Humalog and Regular insulins are clear, others are cloudy. Long and Intermediate acting insulins must be gently mixed by rolling the vial between palms. There should be no clumping of particulate in the insulin. Do not use insulin that is not uniform in consistency.

Steps for Insulin Injection

- 1. Get supplies.
- 2. Wash hands.
- 3. Roll bottle to mix. Wipe top with alcohol swab.
- 4. Pull plunger down to _____
- 5. Push needle into bottle. Push plunger up.
- 6. Pull plunger down to _____ units.
- 7. Pick injection site. Wipe with alcohol swab.

<u>Select injection site.</u> Injections may be given in the abdomen, thighs, buttocks or arms. Insulin sites should be rotated in order to avoid tissue damage, which results in the poor absorption of the insulin. Speed of absorption decreases with each of the following sites: arms, legs and buttocks.

Care and Storage

Effectiveness of insulin is dependent on its careful handling and storage. Date the insulin when it is opened and discard 30 days after opening. Check the expiration date on stored insulin regularly.

- Keep insulin refrigerated for longer shelf life, if a refrigerator is not available, a cool pack may be used. Unrefrigerated insulin should be kept as cool as possible.
- Do not let insulin freeze, if it does, discard it immediately.
- Keep insulin away from heat and light.
- Clumping or frosting results from too much shaking or rough handling. Discard.
- Insulin may be carried in a fanny pack or backpack with an ice pack, as long as
 it is positioned so it does not freeze or get too warm.

ROLES AND RESPONSIBILITIES

The well being of a child with diabetes requires a collaborative relationship between the school and home. The child and his/her family and the health care team are responsible for overall care planning and management. The school is responsible for assuring that the care plan is implemented and supported in the school setting, and that all factors related to the child's diabetes care are communicated to the family.

The staff section of the manual is designed to be individualized for each child and copied for each team member.

Parent/Care Giver/Guardian

- Advocate on behalf of the child
- Participate in the parent conference and planning meetings with school personnel
- Approve the care plan and emergency procedures
- Provide written permission for sharing of medical information with school nurse
- Provide medical equipment, prescriptions, supplies, etc. as indicated in the plan
- Provide food packs for several locations in school and on the bus
- Keep the school informed of any changes in plan of care
- Assist with the staff training if desired

Student

- Consistent with their ability, willingness and parental guidance, students may help:
- Monitoring and recording her/his blood glucose
- Informing adults of symptoms of potential emergencies
- Eating the right foods, in the right amounts, on time
- Exercising regularly
- Carrying supplies for possible hypoglycemic reactions

The Healthcare Team

The child's health care team works with the family to educate them about diabetes and day to day management. They have helped the family and child to make the needed life-style changes and developed the overall plan of care. The child may see one or more members of the team as often as every three months. It is critical that the services provided in school support and assist in the implementation of the overall care plan. Members of the child's health care team may include: pediatric endocrinologist, pediatrician or family practitioner, nurse practitioner, certified diabetes educator (CDE), dietitian (RD).

Principal

General Role:

- Learn about diabetes
- Promote a supportive learning environment
- Treat the child normally and help other children do the same
- Participate in development of the individual care plan
- Understand one's own role and the role of others
- Know what to do in an emergency and the order of responsibility for emergency care
- Know the phone numbers of the child's parent(s and physician

Specific Responsibilities:

- Support the school nurse and care team in the implementation of the individualized care plan throughout the school
- Know regulations that apply to children with diabetes and assure staff compliance
- Explain laws and regulations to community members if necessary
- Facilitate problem solving and negotiations among members of the school team
- Designate and coordinate the 504 team
- Delegate nursing duties if nursing staff is not full time
- Assure substitute teachers know that the child has diabetes and that they are familiar with classroom procedures

Care Plan Implementation Role/Notes From Planning Meeting

Child's name Planned snack times
Usual signs/symptoms of low blood glucose:
Call order for assistance (name and extension):
Other notes:

School Nurse

General Role:

- · Learn about diabetes
- Promote a supportive learning environment
- Treat the child normally and help other children to do the same
- Participate in development of the individual care plan
- Understand one's own role and the role of others
- Know what to do in an emergency and the order of responsibility for emergency care
- Know the phone numbers of the child's parent(s and physician

Specific Responsibilities:

- Generate a nursing assessment of the child based on home or school visit
- Obtain pertinent medical and psychosocial information
- Assure the participation of the parents and student in development of the Individual Care Plan, Emergency Plan, staff training and other aspects of care management in the school
- Coordinate the student's in-school health care as specified in the Health Care Plan
- Organize and conduct Pre-planning and Planning Meetings
- Ensure that caregivers in the school have received competency-based training in child specific techniques and problem management
- Communicate with school team members on a regular basis
- Maintain appropriate documentation of care provided
- Regularly review and update the Individual Care plan and training of caregivers
- · Serve as child advocate

Care Plan Implementation Role/Notes From Planning Meeting

Child's name Planned snack times

Usual signs/symptoms of low blood glucose:

Call order for assistance (name and extension):

Other notes:

Classroom and Substitute Teacher

General Role:

- Learn about diabetes
- Promote a supportive learning environment
- Treat the child normally and help other children to do the same
- Participate in development of the individual care plan
- Understand one's own role and the role of others
- Know what to do in an emergency and the order of responsibility for emergency care
- Know the phone numbers of the child's parents and physician

Specific Responsibilities:

- Learn to recognize the signs and symptoms of hypo- and hyper-glycemia and be able to respond in accordance with the emergency plan
- Help the child to comply with food and snack requirements and monitoring routines
- Communicate diabetes-related needs to substitute and special teachers and instructional assistants
- Educate the class about the special needs of the child (with his/her permission)
 with diabetes and use this as an opportunity to educate students regarding diet,
 exercise and health. The student with diabetes may be willing to present
 information to his or her classroom peers that can enhance peer knowledge
 and support for the student
- Guide the student in choosing a buddy who will be an advocate for him/her in the event of an emergency
- Assure substitute teachers know that the child is diabetic and that they are familiar with classroom procedures

Processing the second
Care Plan Implementation Role/Notes From Planning Meeting
Child's name Planned snack times
Usual signs/symptoms of low blood glucose:
Call order for assistance (name and extension):
Other notes:

Food Service Director and Staff

General Role:

- Learn about diabetes
- Promote a supportive learning environment
- Treat the child normally and help other children to do the same
- Participate in development of the individual care plan
- Understand one's own role and the role of others
- Know what to do in an emergency and the order of responsibility for emergency care
- Know the phone numbers of the child's parent(s)s and physician

Specific Responsibilities:

- Guide the meal modification process
- Assure that food is prepared and served according to the care plan
- Collaborate with the family and child on a menu plan when school meals are chosen
- Work with the school nurse to obtain a medical statement for meal modification
- Keep medical statements on file and update as necessary
- Communicate with the school's nutrition consultant when necessary to implement complex recommendations
- Train the food service staff on meal modification and the needs of the child

Note: The responsibility of the Food Service is to accommodate the medical needs of the child, not personal food preferences. When menus are reviewed and appropriate substitutions are offered, the family may choose to occasionally pack a child's lunch.

Care Plan Implementation Role/Notes From Planning Meeting

Usual signs/symptoms of low blood glucose:	

Call order for assistance (name and extension):

Other notes:

Child's name

Planned snack times

Physical Education Teacher and Coach

General Role:

- Learn about diabetes
- Promote a supportive learning environment
- Treat the child normally and help other children to do the same
- Participate in development of the individual care plan
- Understand one's own role and the role of others
- Know what to do in an emergency and the order of responsibility for emergency care
- Know the phone numbers of the child's parent(s and physician

Specific Responsibilities:

- Be sure blood glucose level is checked before and/or activity if this is in the care plan
- Be aware of the student's usual level of activity
- Allow snacks before or after physical activity if indicated
- Encourage increased fluid consumption
- Learn to recognize the signs and symptoms of hypo- and hyper-glycemia and be able to respond in accordance with the emergency plan
- Communicate diabetes-related needs to substitute and special teachers and instructional assistants
- Encourage participation in sports and athletic opportunities as appropriate
- Ensure that glucose monitoring equipment and "low-packs" are available at all activity sites

Care Plan Implementation Role/Notes From Planning Meeting
Child's name Planned snack times
Usual signs/symptoms of low blood glucose:
Call order for assistance (name and extension):
Other notes:

Counselor

General Role:

- Learn about diabetes
- Promote a supportive learning environment
- Treat the child normally and help other children to do the same
- Participate in development of the individual care plan
- Understand one's own role and the role of others
- Know what to do in an emergency and the order of responsibility for emergency care
- Know the phone numbers of the child's parents and physician

Specific Responsibilities:

- Assist the student with concerns he or she may have regarding their diabetes
- Communicate with the child, family and school staff as necessary
- Respond to ineffective coping mechanisms demonstrated by child and family
- Support the child, family and school personnel in compliance with the treatment plan
- Coordinate a class discussion about diabetes and management needs (if the child doesn't mind)
- Encourage participation in self-management that is appropriate to the child's developmental level

Care Plan Implementation Role/Notes From Planning Meeting

Child's name Planned snack times	
Usual signs/symptoms of low blood glucose:	
Call order for assistance (name and extension):	
Other notes:	

Bus Driver

General Role:

- Learn about diabetes
- Promote a supportive learning environment
- Treat the child normally and help other children to do the same
- Participate in development of the individual care plan
- Understand one's own role and the role of others
- Know what to do in an emergency and the order of responsibility for emergency care
- Know the phone numbers of the child's parents and physician

Specific Responsibilities:

- Learn to recognize the signs and symptoms of hypo- and hyper-glycemia and be able to respond in accordance with the emergency plan
- Know that the end of the school day is often the time of low blood glucose episodes
- Communicate diabetes-related needs to substitute drivers and transportation assistants
- Allow the child to consume a snack on the bus as indicated in their plan.
- Keep a "low-pack" food kit, provided by the family, readily available on the bus

Care Plan Implementation Role/Notes From Planning Meeting

Child's name Planned snack times	
Usual signs/symptoms of low blood glucose:	
Call order for assistance (name and extension):	
Other notes:	

Comparison of Manifestations of Hypoglycemia and Hyperglycemia

Characteristic	Hypoglycemia	Hyperglycemia
Onset	Rapid (minutes)	Gradual (days)
Mood	Labile, irritable, nervous,	Lethargic
	weepy, combative	
Mental Status	Difficulty concentrating,	Dulled sensorium, confused
	speaking, focusing,	, , , , , , , , , , , , , , , , , , , ,
	coordinating	
Inward feeling	Shaking feeling, hunger,	Thirst, weakness, nausea/
	headache, dizziness	vomiting, abdominal pain
Skin	Pallor, sweating	Flushed, signs of
		dehydration
Mucus membranes	Normal	Dry, crusty
Respiration's	Shallow	Deep, rapid (Kussmaul)
Pulse	Tachycardia	Less rapid, weak
Breath odor	Normal	Fruity, acetone
Neurologic	Tremors Late:	Paresthesia
	hyperreflexia, dilated	
	pupils, convulsion	
Ominous signs	Shock, coma	Acidosis, coma
Blood		
Glucose	Low: below 70 mg/dl	High: 240 mg/dl or more
Ketones	Negative/ trace	High/large (only if DKA)
Osmolarity	Normal	High
pH	Normal	Low (7.25 or less)
Hematocrit	Normal	High
НСоз	Normal	Less than 15eEq/L
Urine:		
Output	Normal	Polyuria to oliguria
Sugar	Negative	High
Acetone	Negative/trace	High

AUTHORIZATION FOR RELEASE OF MEDICAL INFORMATION

To:	
To:(Doctor's Name)	
From: (Your name and address)	
Please send information about my child Date of birth:/ To:	
Information to be sent:	_ _
Problem list only	
An update on	
Ongoing progress report on	
(O' (D (10 1)	

(Signature of Parent/Guardian)

Prescription medication order and permission form

(to be returned to the school nurse)

From time to time, it may be necessary for your child to take prescription medicine for treatment of an illness. Medicines that are ordered to be taken less than 4 times a day can and should be taken at home. However, if medicine must be taken 4 times a day, or at a specific time scheduled during school hours, the school nurse, as mandated by state law, may dispense medications **ONLY WITH THE FOLLOWING:**

- 1. Medication order, signed by the physician
- 1. Parental authorization, signed by the parent
- 2. Original pharmacist labeled bottle

	Medication Order
Student:	
Medication:	
Reason for giving:	
	phone number of physician:
Signature of Physician)	
	PERMISSION TO ADMINISTER
Date:/	
I hereby give my permission	n for
to take the above prescription	on at school as directed.
(Signature of Parent/Guar	dian)

APPENDIX 4 Medical Statement for Children Requiring Modifications in School Meals

Name of Student:

Name of Parent/Guardian:		Daytime Phone:		
L				
Disability or Medical Condition requiring modification of school meals:	Major life activity affected by student's disability (please circle all that apply): caring for one's self, eating, performing manual tasks, walking, seeing, hearing, speaking, breathing, learning, working.			
Required Meal Modification (check	(all which apply):			
RESTRICTED NUTRIENT	INCREAS	ED NUTRIENT	MODIFIED TEXTURE	
Calorie Controlled Carbohydrate Protein Sodium Fat/Cholesterol	Calorie Protein Fiber Other:		Describe required modification:	
FOODS TO BE OMITTED FI	ROM THE DIET			
List all that apply:		Foods that may b	e substituted:	
Special Utensils Needed:				
Tube Feeding Required:				
Other Accommodations needed:				
For student with a disability:	Signature	of Physician:		
	Date:			
For non-disabled student:	Signature	of Other Medical A	uthority:	
RECOMMENDATIONS FOR MANAGEN	Date:	WITH DIABETES IN	SCHOOL	

Birth date:

Sample--504 Plan for the School Year 97/98

Student Name: <u>Mary Jones</u>	_Date	e of Birth: 11/15/ 89	Grade: <u>4</u>
Date of Disability Determination: 8 / 15 /	98	Case Manager: Susar	Doe, RN

Date of Rirth: 11/15/ 90

Area of disability that "substantially limits a major life activity"/Description of how this disability limits a major life activity:

- Mary has type I diabetes mellitus. This is a condition in which the pancreas is unable to make insulin. Without insulin, the body cannot change glucose (sugar) into the energy a person needs. To compensate for the lack of natural insulin, she must take daily insulin injections, usually at home but sometimes in school.
- Mary's daily insulin injections must be balanced with her meals, snacks and regular physical activity. To consistently achieve this balance, she must eat daily snacks and meals on a regular schedule. During the school day she must check her blood glucose before lunch, and physical education class, as well as when her body tells her blood glucose is low or too high.
- While Mary is achieving independence in self-management of her diabetes, the adults who work with her will need to be supportive and understanding about the daily regimen. Her self-care needs will be integrated into the school day so there are minimal interruptions in the learning environment.
- Mary is generally responsible and independent about her blood glucose tests, diet, and necessary equipment. The adults in the school community will help by reminding Mary to bring her pack on all out-of-school trips and to keep it with her when she is away from the classroom for extended periods of time. Mary will need this reminder especially before special events.
- Mary's blood glucose levels affect the way he/she learns.
- Mary's behavior is related to blood glucose levels. She can feel "racey" and excited when her blood glucose is high, tired and "spacey" when it is low.
- When Mary is excited and/or stressed as in a testing situation, her blood glucose can potentially go up. When her blood glucose is high (over 200) her body responds by trying to decrease this glucose level. She may become thirstier as her body is acting to dilute or flush out the extra sugar. She needs to drink more water and then urinate more frequently.
- The learning environment is altered when Mary must stop an activity to test her blood sugar, go to the bathroom, eat a snack or get a drink of water.
- Mary must continuously remind herself to monitor her blood sugar at appropriate times, to eat/exercise regularly and to bring supplies with her. This self-monitoring is a big task and is a distraction in itself.

RECOMMENDATIONS FOR MANAGEMENT OF CHILDREN WITH DIABETES IN SCHOOL

Home/School Communication: To develop parent/student school communications:

- 1. There will be on going communication between parents and case manager.
- 2. Parent-teacher meetings will be scheduled at regular times especially at the beginning of the school year and other transition times. Parents want to be contacted immediately if any academic or social concerns arise.
- 3. Health concerns will be addressed as the need indicates.
- 4. Consistency is important in Mary's academic plan. Transition meetings including last and current teaching staff, nursing and parents will be scheduled.
- II **Organization/Management:** To modify the instructional day:
 - 1. Mary will need modification of non-academic time (long lunch, extra snack period on occasion).

III Alternative Teaching Strategies/Accommodations: To modify teaching methods:

- 1. Adjust testing procedures:
 - Mary may need to have open bathroom privileges during standardized tests.
 She should be seated so she can come and go from the room easily. Please remind her that she can go at anytime despite testing rules.
 - Mary must keep her low pack with equipment and snacks next to her in the testing area.
 - If Mary goes to the bathroom before or during the testing, special accommodations may need to be made to repeat instructions or to lengthen her testing time.
 - Mary may need to have snack at different times and intervals than the other students.
- 2. Individualize classroom/homework assignments:
 - There will be a need to explain assignments or adjust them if Mary's blood glucose is unusually high or low on certain days. Her level of concentration is affected and she will need accommodations.
- 3. Repeat or structure instructions for in-class or homework assignments.
 - Mary may miss part of an assignment or a class while testing, in the bathroom or eating his/her snack. Repeating verbal/written instructions will be needed.
- If Mary's blood glucose is unusually high or low (>300 or < 80), she may feel shaky, slightly disoriented or very distracted. The classroom teacher will need to revisit instructions or concepts she may have missed in class.

IV <u>Student Precautions:</u> Please see attached Diabetic Day Plan Evaluation data used to make this decision:

- Diagnosis of Diabetes Mellitus Type I
- Parents Report and past concerns
- Teachers observation in class
- Academic performance
- Articles and data provided by parents

The following related aids and services are recommended:

- 1. Health services: The school nurse will be available on a daily basis to provide support and guidance to Mary and the school staff. A trained substitute nurse will be available if Susan Doe is not in school.
 - Mary's classroom teacher and one other adult staff member will be trained in the administration of Glucagon.
 - Susan Doe, RN, will inform all appropriate teaching staff and cafeteria staff about condition and what to do in an emergency.
- 2. Equipment and Food Items
 - Mary will carry her own glucometer, finger lancets and glucose strips. She will also carry a "low pack" with juice, and a snack.
 - There will be extra juice, and snacks kept in the classroom area, the library and in exploratory classes as needed.
 - The nurse's office will have extra juice, crackers, peanut butter and other snacks for Mary and will keep a vial of insulin, extra glucose strips, syringes and ketone strips for urgent use. A Glucagon kit will be kept in the locked medicine cabinet in the nurse's office.
 - Mary's parents will provide all food and equipment to the school. Susan Doe, RN, will be responsible for distributing the food and maintaining the supply.

Sample Student Accommodation Plan

Studer	nt Name: <u>Mary Jones</u>	_Date of Birth: 11/15/ 89 Grade: 4
Case I	Manager: <u>Susan Doe, RN</u>	_School: Central
Date c	of Meeting: <u>8 / 15 / 98</u>	
1.	Describe the nature of the conce Type 1 diabetes	rn:
2.	Describe the basis for the determ Type 1 diabetes is a physiological	nination of disability: al disorder that affects the endocrine system.
3.	Describe how the disability affect The student is at risk for hype an metabolic dysfunction.	s a major life activity: d hyperglycemic episodes related to her/his
4.	 Treat or receive assistance w or Glucagon as prescribed) Treat or receive assistance w insulin if ordered and drink was Eat whenever and wherever reflected and unrestricted acceptable. 	e with blood glucose monitoring ith treatment of hypoglycemic episodes (glucose and ith treatment of hyperglycemic episodes (inject ater)
Revie	w/Reassessment Date://_	<u>—</u>
Partici	pants (name and title):	

• Attachment: Information regarding Section 504

CC: Student's Cumulative File

Universal Precautions and Infection Control

Schools should have established procedures for responding to universal precautions and infection control. The following is provided as general guidance.

In response to the increase in hepatitis B and human immunodeficiency virus (HIV) infections, the Centers for Disease Control have recommended "universal blood and body-fluid precautions." These measures are intended to prevent transmission of these and other infections, as well as to decrease the risk of exposure for care-providers and students. As it is currently not possible to identify all infected individuals, these precautions must be used with every student, regardless of their medical diagnosis.

Universal precautions pertain to blood and body fluids containing blood, cerebrospinal fluid, synovial fluid, vaginal secretions, semen, and pericardial fluid. These precautions do not apply to other body products such as saliva, sputum, feces, tears, nasal secretions, vomitus and urine unless blood is visible in the materials. However, these other fluids and body wastes can be sources of other infections and should be handled as if they are infectious.

The single most important step in preventing exposure to and transmission of any infection is anticipating potential contact with infectious materials in routine as well as emergency situations. Based on the type of possible contact, the caregiver should be prepared to use the appropriate precautions and techniques prior to providing care. Diligent and proper hand washing, the use of barriers, appropriate disposal of waste products and needles, and proper decontamination of spills are essential techniques of infection control. Using common sense in the application of these measures will enhance protection of both the caregiver and the student.

Hand Washing

Proper hand washing is crucial to preventing the spread of infection. Textured jewelry on the hands or wrists (such as rings with stones) should be removed prior to washing and kept off until completion of the care procedure and hands are rehashed. Use of running water, lathering with soap and using friction to clean all surfaces of remaining jewelry and hands is key. Rinse well with running water and dry hands with paper towels. If soap and water are unavailable, wet towelettes or "Handi-wipes" may be used.

- Hands should be washed before physical contact with student and after the contact is completed.
- Hands should be washed after contact with any used equipment.

- If hands (or other skin) become soiled with blood or body fluids, they should be washed immediately before touching anything else.
- Hands should be washed whether gloves are worn or not and after gloves are removed.

Barriers

Barriers include disposable gloves, protective eyewear, masks and gowns. The use of barrier is intended to reduce the risk of contact with blood and body fluids for the caregiver as well as to control the spread of infectious agents from student to student. It is essential that appropriate barriers be used when contact with potentially infectious materials is possible.

Gloves should be worn when direct care of the student may involve contact with blood or body fluids. For infection control, it is recommended that gloves be worn as well for contact with urine, feces and respiratory secretions. Gloves should be disposed of after each use and not reused.

Disposal of Waste

Needles, syringes and other sharp objects should be placed in a metal or other punctured-proof container immediately after use. To reduce the risk of an accidental needle stick or cut, needles should not be recapped, bent or removed from the syringe before disposal. Once it is full, the container should be sealed, double bagged and then disposed of in the garbage away from the reach of children.

INSTRUCTIONS FOR COMPLETING THE MEDICAL STATEMENT

This form may be completed by a parent but must be signed by a physician if the child has a disability. A Major Life Activity must be circled if the child has a disability. In many cases it is "eating."

If the child does not have a disability it may be signed by a recognized "medical authority" which includes a physician, physician assistant, registered dietitian, registered nurse, or occupational therapist or other health professional specified by the Nevada Department of Education.

Check the required meal modification(s) the student needs. Both the modification category and the detailed type of modification should be checked. The more information provided, the better able the school is to meet the student's needs.

Food omission is most often needed due to a food allergy. Specify to what extent a food must be avoided. For example: "omit milk as a beverage", "omit foods which have milk or other dairy products as a major ingredient," "milk and all dairy products must be completely omitted from the diet."

Food to be substituted: Be as specific as is reasonable. Typical substitutions would be: "juice for milk," "any other vegetable for tomatoes," "equivalent menu item which does not contain eggs," "fresh or unsweetened fruit for dessert," etc.

Special Utensils refers to special silverware, plates, cups or other items the student needs in order to eat the meal.

Other information includes needs which do not directly relate to the modification of a food such as: fluid intake at other than meal time; additional time to eat or specific timing of a meal or snack; feeding techniques.

Note: The responsibility of the Food Service is to accommodate the medical needs of the child, not personal food preferences. When menus are reviewed and appropriate substitutions are offered, the family may choose to occasionally pack a child's lunch.

Once completed, this form should be returned to the Food Service Manager who, with the input of the parents and appropriate school staff, will establish the necessary routines to provide the modified meal. The original copy of this Medical Statement should be kept on permanent file in the nurse's office or food service office. It remains in effect until replaced or inactivated.

Data/Information Sheet—Diabetes Care in School

To be completed by parent or legal guardian

Child's Name			[Date of Birth	
Mother's Name		Fathe	er's Name		
Child's age at time of	of diagnosis _				
Contact phone numbers parent, friend, or car person has legal au	regiver and wh	nether the nu	mber is home	e and or work. Als	
		Call Order			
Number	Name	Relationship	Location	Authority	
Example 333-5442					
1.					
2. 3.					
Δ					
5.					
Primary Care Physic	cian·				
Name:			phone:		
Other Health Provid	ers:				
Name:			phone:		_
Name:					_
Name:			priorie		_
Monitoring: Usual E	Blood glucose	monitoring til	mes:		
Morning	-		Night		
Mid-day			Other		
Afternoon			Other		
Does your child hav	e a preferred l	location for m	onitoring (cla	assroom, health of	ffice, etc.)?

47

RECOMMENDATIONS FOR MANAGEMENT OF CHILDREN WITH DIABETES IN SCHOOL

Data/Information Sheet—page 2. FOR (Child's name)		
Goal range for blood glucose: Between and	·	
Lows: If below, please do the following: Call parent if below:		
Usual signs/symptoms of low blood glucose are:		
Highs: If above, please do the following:		
Blood glucose level at which to measure ketones:		
Call parent if above:		
Usual signs/symptoms of high blood glucose are:		
Insulin (circle the answer's below): Will daily insulin be needed in school?	YES	NO
Will insulin be needed in school at any other time? If YES to either, identify time, amount and circumstances for a insulin.	_	NO
Food: Will child participate in school breakfast and/or school lunch?	YES	NO
If yes, will modifications to the regular menu be needed?	YES	NO
What are usual times for meals/snacks?		
Mid-morning	Mid-afternoo Supper Bed-time	n

RECOMMENDATIONS FOR MANAGEMENT OF CHILDREN WITH	DIABETES IN SCHOOL	
Data/Information Sheet—page 3. FOR (Child's name)		_
Exercise: What are your child's favorite physical activities?		
Will your child participate in school sports?	YES NO	

Self Care:

Please put an X in the box that best describes your child's role:

	Does it	Does it with	Helps	Parent
	alone	supervision	parent	does it
Picks finger				
Puts strip in monitor				
Reads monitor				
Records result				
Adjusts food based on result				
Adjust insulin based on result				
Knows which foods to limit				
Can select kind and amount of				
food				
Helps plan meals				
Selects insulin injection site				
Prepares for injection, selects				
site				
Injects insulin				
Measures insulin				
Determines amount and type of				
insulin				
Measures ketones				
(Other)				

RECOMMENDATIONS FOR MANAGEMENT OF CHILDREN WITH DIABE	TES IN SCHO	JOL	
Data/Information Sheet—page 4. FOR (Child's name)			
Parties and Special Food or Meals:			
Do you wish to be contacted before each event?	YES	NO	
Additional instructions for the school:			
Emergencies:			
What do you feel should be treated as an emergency?			
What do you want the school to do in an emergency?			

Other Requests:

Parent Checklist

This checklist is provided to help parents identify the forms, supplies and other materials they need to bring to the school. The list needs to be modified for individual children.

All iter	ms on the checklist should be sent to the school nurse.
	Data / Information form
	Photograph of child
	Signed Release of Information for physician(s)
	Monitoring supplies: Lancets, meter, strips, alcohol, ketone strips, etc.
	Snack (Low) packs. Number:
	Glucose tablets, Gel (tubes). Number
	Record keeping sheets
	Insulin and related supplies: Syringes, alcohol, etc.
	Prescription Medication Order and Permission form for Insulin
	Glucagon kits with premeasured dosage. Number
	Prescription Medication Order and Permission form for Glucagon

Individual Care plan— Diabetes Care in School

	Child's Name : Grade:
Picture Here	

Contact phone numbers (in priority order):

Call	Number	Name, relationship	Location
1.			
2.			
3.			
4.			
5.			

Primary	Care Physician:		
Name: _		_ phone:	

Routine blood alucose monitoring

Time Place Who			
1.			
2.			

Indications for special blood glucose monitoring:

Individual Care Plan--page 2.

Usual signs/symptoms of Low blood glucose:

High blood glucose:

Actions required by blood glucose levels in mg/dl- (call parents/guardians, give snacks, measure ketones, etc.):

	Glucose Level	Action
1.		Normal No Action needed
2.		
2.		
4.		
5.		

Food

	Time	What (snack, lunch. other)
1.		
2.		
3		

Requires menu modification by Food Service:

YES NO

If yes, form has been completed and is on file in Food Service Office: YES NO

Parties, Special Occasions—(Special considerations, if indicated by parents)

Physical activity/exercise/ sports—(Special notes, if indicated by parents)

Individual Care Plan--page 3.

Insulin-time or circumstance (e.g. Blood Glucose over 300mg/dl) when to administer and who will be responsible:

	Time or ?	Place	Who
1.			
2.			

Glucagon:

Will school staff administer Glucagon?

YES NO

If yes, describe circumstances when it should be given.

Does the child weigh less than 45 pounds? If yes, note on Glucagon kit. YES NO Location of supplies in the school:

Site	Staff Name	Extension	Notes
Monitoring for blood glucose			
Snack or low packs	, glucose tablets or gel		
Snack or low packs	, glucose tablets or gel		
1	, <u>J</u>		

RECOMMENDATIONS FOR MANAGEMENT OF CHILDREN WITH DIABETES IN SCHOOL

Individual Care Plan—page 4. Disposal of Supplies:					
Emergency Plan: Situation(s) constituting and emergency:					
Action Needed:					
Phone Numbers:					
Number	Who				
	Rescue Squad				
	Parent				
Goals: Describe goals for child 's care in the sch 1.	nool setting for the year.				
2.					
3.					
Signatures:					
Parent	School Nurse				
Principal					

Individual Care Plan—page 5.

Staff Training Record

		T : D	ing Record	
	Date Training Received			
Staff Member	Diabetes	Monitoring	Insulin	Notes
Name	Basics		and	
	Bacico		Glucagon	
			Glucagon	
	1			
-				

SUPPLEMENTAL INFORMATION SHEET FOR STUDENT WEARING AN INSULIN PUMP AT SCHOOL (LICENCED HEATHCARE PROVIDER'S ORDERS FOR INSULIN PUMPS FOR HEATH SERVICES AT SCHOOL)

School YearStudent Name: Date of Birth: Pump Brand/Model:
Pump Resource Person: Phone/Beeper (See diabetes care plan for parent phone#)
Blood Glucose Target Range: - Pump Insulin: Humalog □ Regular □
Insulin Correction Factor for Blood Glucose Over Target:
Insulin: Carbohydrate Ratios:
(Student to receive insulin bolus for carbohydrate intake <i>immediately before</i> / <i>minutes</i> before
eating. Circle appropriate interval)
Location of Extra Pump supplies
INDEPENDENT MANAGEMENT This student has been trained to independently perform routine numb management and troublesheet
This student has been trained to independently perform routine pump management and troubleshoot problems, including but not limited to:
 Giving boluses of insulin for both correction of blood glucose above target range and for food
consumption.
Changing of insulin infusion sets using universal precautions
 Switching to injections should there be a pump malfunction. Parents will provide extra supplies to include infusion sets, reservoirs, batteries, pump insulin,
syringes.
Symiges.
NON-INDEPENDENT MANAGEMENT (Child Lock On? Yes No circle one) Because of young age or other factors, this student cannot independently evaluate pump function nor independently change infusion sets. Insulin for meals and snacks will be given and verified as follows:
Insulin for correction of blood glucose over will be given and verified as follows:
PARENT NOTIFICATION: Refer to basic diabetes care plan and check all others that apply. Contact
parent in event of:
Pump alarms/malfunction
Soreness or redness at site
Detachment of dressing/infusion set out of place
Leakage of insulin
☐ Student must give insulin injection ☐ Student has to change site
Corrective measures do not return blood glucose to target range within hrs.
Other:

MANAGEMENT OF HIGH/VERY HIGH BLOOD GLUCOSE: Refer to previous sections and to basic Diabetes Care Plan

MANAGEMENT OF LOW BLOOD GLUCOSE Follow instructions in basic Diabetes Care Plan, but in addition:

If low blood glucose recurs without explanation, notify parent/diabetes provider for potential instructions to suspend pump.

If seizure or unresponsiveness occurs:

Give Glucagon and/or glucose gel (See basic Diabetes Health Plan)
 Call 911
 Notify parent
 Stop insulin pump by:

 Placing in "suspend" or stop mode
 Disconnecting at pigtail or clip

Cutting tubing 5. If pump was removed, send with EMS to hospital. COMMENTS:	
Effective Dates: From: To:	
Parent's Signature:	Date:
School Nurse's Signature:	Date:
Diabetes Care Provider Signature:	Date: